

# Health Consultation

---

**PUBLIC COMMENT RELEASE**

SWANN PARK  
296 WEST MCCOMAS STREET

BALTIMORE, MARYLAND

JUNE 13, 2007

**COMMENT PERIOD END DATE: JULY 30, 2007**

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
Public Health Service  
Agency for Toxic Substances and Disease Registry  
Division of Health Assessment and Consultation  
Atlanta, Georgia 30333

## **Health Consultation: A Note of Explanation**

An ATSDR health consultation is a verbal or written response from ATSDR to a specific request for information about health risks related to a specific site, a chemical release, or the presence of hazardous material. In order to prevent or mitigate exposures, a consultation may lead to specific actions, such as restricting use of or replacing water supplies; intensifying environmental sampling; restricting site access; or removing the contaminated material.

In addition, consultations may recommend additional public health actions, such as conducting health surveillance activities to evaluate exposure or trends in adverse health outcomes; conducting biological indicators of exposure studies to assess exposure; and providing health education for health care providers and community members. This concludes the health consultation process for this site, unless additional information is obtained by ATSDR which, in the Agency's opinion, indicates a need to revise or append the conclusions previously issued.

You May Contact ATSDR Toll Free at  
1-800-CDC-INFO

or

Visit our Home Page at: <http://www.atsdr.cdc.gov>

HEALTH CONSULTATION

*PUBLIC COMMENT RELEASE*

SWANN PARK  
296 WEST MCCOMAS STREET

BALTIMORE, MARYLAND

Prepared By:

Division of Health Assessment and Consultation and  
Division of Regional Operations  
Agency for Toxic Substances and Disease Registry

*This information is distributed solely for the purpose of pre-dissemination public comment under applicable information quality guidelines. It has not been formally disseminated by the Agency for Toxic Substances and Disease Registry. It does not represent and should not be construed to represent any agency determination or policy.*

## Table of Contents

Foreword.....	iii
Summary.....	1
Statement of Issues and Community Health Concerns.....	2
Statement of Issues.....	2
Swann Park Community Concerns.....	3
Background.....	4
Site Description and History.....	4
The Swann Park Community:.....	5
Results.....	7
Public Health Implications.....	13
Conclusions, Recommendations, and Public Health Action Plan.....	19
Authors.....	20
References.....	21
Attachment 1.....	23
Attachment 2.....	27

## List of Figures

Figure 1. Location of Swann Park and surrounding community.....	6
---	---

## List of Tables

Table 1. Summary of arsenic and kepone surface soil concentrations from Swann Park.....	8
Table 2. Exposure scenarios and assumptions for people using Swann Park.....	9
Table 3. Soil adherence rates for kepone dermal dose estimation.....	10
Table 4. Calculated doses for arsenic and kepone at Swann Park; 2007 soil data.....	12
Table 5. Community concerns related to Swann Park and ATSDR responses.....	17

## FOREWORD

The Agency for Toxic Substances and Disease Registry, ATSDR, was established by Congress in 1980 under the Comprehensive Environmental Response, Compensation, and Liability Act, also known as the *Superfund* law. This law set up a fund to identify and clean up our country's hazardous waste sites. The Environmental Protection Agency, EPA, and the individual states regulate the investigation and clean up of the sites.

Since 1986, ATSDR has been required by law to conduct a public health assessment at each of the sites on the EPA National Priorities List. The aim of these evaluations is to find out if people are being exposed to hazardous substances and, if so, whether that exposure is harmful and should be stopped or reduced. If appropriate, ATSDR also conducts public health assessments when petitioned by concerned individuals. Public health assessments are carried out by environmental and health scientists from ATSDR and from the states with which ATSDR has cooperative agreements. The public health assessment program allows the scientists flexibility in the format or structure of their response to the public health issues at hazardous waste sites. For example, a public health assessment could be one document or it could be a compilation of several health consultations - the structure may vary from site to site. Nevertheless, the public health assessment process is not considered complete until the public health issues at the site are addressed.

**Exposure:** As the first step in the evaluation, ATSDR scientists review environmental data to see how much contamination is at a site, where it is, and how people might come into contact with it. Generally, ATSDR does not collect its own environmental sampling data but reviews information provided by EPA, other government agencies, businesses, and the public. When there is not enough environmental information available, the report will indicate what further sampling data is needed.

**Health Effects:** If the review of the environmental data shows that people have or could come into contact with hazardous substances, ATSDR scientists evaluate whether or not these contacts may result in harmful effects. ATSDR recognizes that children, because of their play activities and their growing bodies, may be more vulnerable to these effects. As a policy, unless data are available to suggest otherwise, ATSDR considers children to be more sensitive and vulnerable to hazardous substances. Thus, the health impact to the children is considered first when evaluating the health threat to a community. The health impacts to other high risk groups within the community (such as the elderly, chronically ill, and people engaging in high risk practices) also receive special attention during the evaluation.

ATSDR uses existing scientific information, which can include the results of medical, toxicologic and epidemiologic studies and the data collected in disease registries, to determine the health effects that may result from exposures. The science of environmental health is still developing, and sometimes scientific information on the health effects of certain substances is not available. When this is so, the report will suggest what further public health actions are needed.

**Conclusions:** The report presents conclusions about the public health threat, if any, posed by a site. When health threats have been determined for high risk groups (such as children, elderly, chronically ill, and people engaging in high risk practices), they will be summarized in the conclusion section of the report. Ways to stop or reduce exposure will then be recommended in the public health action plan.

ATSDR is primarily an advisory agency, so usually these reports identify what actions are appropriate to be undertaken by EPA, other responsible parties, or the research or education divisions of ATSDR. However, if there is an urgent health threat, ATSDR can issue a public health advisory warning people of the danger. ATSDR can also authorize health education or pilot studies of health effects, full-scale epidemiology studies, disease registries, surveillance studies or research on specific hazardous substances.

**Community:** ATSDR also needs to learn what people in the area know about the site and what concerns they may have about its impact on their health. Consequently, throughout the evaluation process, ATSDR actively gathers information and comments from the people who live or work near a site, including residents of the area, civic leaders, health professionals and community groups. To ensure that the report responds to the community's health concerns, an early version is also distributed to the public for their comments. All the comments received from the public are responded to in the final version of the report.

**Comments:** If, after reading this report, you have questions or comments, we encourage you to send them to us.

Letters should be addressed as follows:

Agency for Toxic Substances and Disease Registry  
ATTN: Records Center  
1600 Clifton Road, NE (Mail Stop E-60)  
Atlanta, GA 30333

## Summary

Recent soil sampling at Swann Park, 296 West McComas Street, in South Baltimore Maryland have found elevated concentrations of inorganic arsenic in surface and subsurface soil samples collected on April 16, 2007. On April 19, 2007, on the recommendation of the Baltimore City Health Department and the Maryland Department of the Environment, the Baltimore City Recreation and Parks Department closed Swann Park. The City Health Department then requested that the Agency for Toxic Substances and Disease Registry (ATSDR) evaluate potential arsenic exposures to users of the park and provide recommendations to protect public health.

Swann Park is in an industrial area of south Baltimore and is directly adjacent to the former site of the Allied Chemical Corporation, Baltimore Plant (2000 Race Street). Until its recent closure, Swann Park was a highly used recreational facility. Activities at the Park included baseball, softball, football, soccer, and lacrosse, as well as general recreation and visitation by community members. In addition to park visitors, the Park workers were constantly engaged in field maintenance activities. All of these park activities may result in exposures to Park soil.

ATSDR screened all of the contaminants detected in the April 2007 Swann Park soil samples against environmental guidelines. Of the many contaminants analyzed and detected in Swann Park soils only arsenic was detected above its environmental guideline. Normal park activities, such as field sports (baseball, football, etc.), and park maintenance have resulted in exposures to the arsenic-contaminated soil. With the exception of pica children (children who eat soil or non-food items), estimates of arsenic doses based on these specific park activities result in doses that are not likely to produce adverse health effects. A child consuming about a teaspoon of soil may experience temporary nausea, diarrhea, vomiting or stomach cramps. These acute effects are unlikely to continue and other long term adverse health effects, including cancer, are unlikely for all exposures.

Dose estimates based on historic soil sampling present the same conclusions as the recent soil data. That is, with the exception of pica children, estimates of arsenic doses based on the above park activities are not likely to produce adverse health effects. However, there may have been an additional dose component from inhalation of airborne plant emissions when the plant was operating. Consequently, the conclusions of this health consultation are limited to the time period after 1976.

Additional collection and analysis of environmental samples for Swann Park is pending. ATSDR will evaluate these data as they become available and revise the conclusions and recommendations of this health consultation as appropriate.

## Statement of Issues and Community Health Concerns

### Statement of Issues

In response to the community health concerns, on April 18, 2007, the Commissioner of the Baltimore City Health Department first contacted the Agency for Toxic Substances and Disease Registry (ATSDR) to request ATSDR advice and recommendations regarding public health exposures to the contamination in the park. The Health Commissioner formally requested that ATSDR conduct a public health evaluation of arsenic exposures at Swann Park, Baltimore Maryland. The purpose of this document is to present the ATSDR evaluation of April 2007 soil sampling data and likely exposures to Park soil to determine if anyone is likely to get sick from playing or working at Swann Park.

Based on historic and recent soil sample data from Swann Park, inorganic arsenic is the primary contaminant of concern. However, the most recent soil samples from Swann Park also included analyses of other metals, pesticides, and semi-volatile organic compounds (SVOCs). Consequently, this health consultation will also evaluate the distribution and concentrations of those analytes to determine if they are present in Swann Park at concentrations of public health concern.

Community exposures to Swann Park soil are evaluated in a two-step process. The first step is to compare maximum measured analyte concentrations with soil screening levels or *environmental guidelines*.<sup>1</sup> Those analytes detected above their respective screening values are “contaminants of (public health) concern” and require further evaluation. The second evaluation step is to calculate potential exposure doses for each contaminant of concern and compare those doses with *health guidelines*<sup>2</sup>, or doses that may be related to specific diseases or adverse health effects. The potential health affects of those contaminants with doses that exceed their respective health guidelines are discussed relative to the types of exposures that have occurred at Swann Park.

Until its recent closure, Swann Park was a highly used recreational facility. Activities at the Park included baseball, softball, football, soccer, and lacrosse, as well as general recreation and visitation by community members. In addition to park visitors, the Park workers were constantly engaged in field maintenance activities. All of these park activities may result in exposures to Park soil.

The following *Community Health Concerns* section of this document presents the specific concerns or questions that people have raised to ATSDR and the City of Baltimore Health Department about their exposures to Park soil. We have used these specific concerns to define

---

<sup>1</sup> *Environmental Guideline*: Calculated concentration of a substance in air, water, food, or soil that is unlikely to cause harmful (adverse) health effects in exposed people. The comparison value (CV) is used as a screening level during the public health assessment process. Substances found in amounts greater than their CVs might be selected for further evaluation in the public health assessment process.

<sup>2</sup> *Health guideline*: Substance-specific doses derived using toxicologic and epidemiologic data with safety factors to ensure that they are protective of human health. When adequate dose-response data are available, guidelines are developed for ingestion and inhalation exposures. Various guidelines are developed by ATSDR and other Federal and State agencies. Specific guidelines used in this assessment are referenced and defined.

how people may have been exposed to park soils and subsequent sections will provide our evaluation of the potential for those exposures to cause sickness or disease.

The Background Section provides brief descriptions and histories of the Race Street chemical plant and Swann Park and the surrounding neighborhood. The *Results* section presents our evaluation of the specific contaminants present in Swann Park soil and the contaminant doses from specific park activities. The section on *Public Health Implications* presents an evaluation of the types of diseases caused by these contaminants and whether the estimated doses are at levels likely to cause those diseases. This section also includes specific responses to all of the community concerns. Finally, the *Conclusions, Recommendations and Public Health Action Plan* section summarizes our findings on Swann Park and our recommendations and proposed actions for addressing public health issues related to contaminated soil in the park.

### **Swann Park Community Concerns**

ATSDR and the Baltimore City Health Department collected community concerns during a public meeting held at Digital Harbor High School on April 26, 2007. In follow up to this meeting, ATSDR interviewed a neighbor to the site, current and former coaches from local high schools that regularly fielded teams at Swann Park, and City employees who maintained the park.

Current community concerns include:

- Nearby residents are concerned about their unique exposures to contamination at the park, including inhalation of dust that would blow from the park towards their homes. One family is concerned in particular about any relationship between exposure to arsenic-contaminated soils and skin conditions, seizures in a toddler, nervous system/mental health conditions, sleep apnea, diabetes, and a fatal blood clot/aneurysm.
- Many community members raised concerns and questions about medical testing options for individuals who were exposed to the contamination at the park. One concern was raised that there is a need for public health authorities to provide some guidance to parents who have already pursued getting hair and nail testing for their children.
- Some community members were interested in long term follow up for people who were exposed to contamination at the park.
- Many concerns were raised about the exposures to student athletes and their coaches. For example, several times a year, students and their coaches would engage in heavy field maintenance activities (raking, moving dirt/muck around, etc), and this would represent a high level of contact with the soils at the site. Many current and former student athletes would be a member of a sports team that would use the field for their entire four-year high school career; a minority of these student athletes would be members of multiple sports teams and therefore would frequent the fields for more than one playing season. Many student athletes would not only use the fields for their team activities, but also during their off hours for general recreation. Athletic team representatives and community members emphasized that there are no facilities for washing hands at the fields, and many users would eat and drink with dirty hands in the midst of activities at the site. One particular concern is the exposures to baseball catchers, who spend a significant amount of their playing time at ground level and in the dirt.

- Community members were interested in how ingestion of arsenic-contaminated soils affects the human body.
- Community members asked how much soil at the highest levels of arsenic contamination found in surface soils at the park would need to be ingested to produce a toxic effect.

## **Background**

### **Site Description and History**

#### **Race Street Site**

A chronology of historic and recent actions at the Allied Signal Race Street Site is available at the Maryland Department of the Environment web site (MDE, 2007). Various insecticides, herbicides, and other arsenic products were produced or processed at the Race Street site from 1897 until 1976 (Matanoski, et al. 1981). The original facility was torn down and replaced in 1952. The rebuilt plant produced arsenic acid, calcium and lead arsenic, cupric acetoarsenite (Paris green), and sodium arsenite. These products were dried and packaged as solid materials except for Paris green, which was shipped as a liquid (not produced after early 1950s). Other pesticides, such as chlorinated hydrocarbons and organophosphates, were not produced at the Plant, but were processed and packaged at the facility.

Allied Chemical (later Allied-Signal) purchased and operated the Plant in 1955 until it was shut down in 1976. Allied-Signal, was in turn, purchased by Honeywell International in 1999. The City of Baltimore purchased the Race Street Property in 1977 and demolition of the factory buildings occurred throughout 1977. Interstate 95 was built on a portion of the property. MDE has overseen several remedial actions, such as placement and maintenance of a clay cap, to limit exposure and migration of site contaminants. An Administrative Consent Order between MDE, the City of Baltimore, and Honeywell International to complete assessment and cleanup of the Race Street Site is currently pending.

#### **Swann Park**

Swann Park is an approximately 11-acre, rectangular park owned by the City of Baltimore, south of the Federal Hill neighborhood (Figure 1). It borders the Middle Branch of the Patapsco River, and has been park land since at least 1914 (Honeywell, 2007a). Until the park was closed on April 19, 2007, the field was used for recreational activities for team sports (e.g., softball, football). For decades, Swann Park was used by Southern High School (now Digital Harbor High School) football and baseball teams, as well as local sports leagues. The park was temporarily closed in 1976, when a pesticide called kepone was found in the dirt of the ball fields. Following completion of remediation (including removal of contaminated soil, plowing, raking, and re-sodding) a panel of federal, state and local officials allowed the park to be reopened that year (Baltimore Sun, 2007).

**Swann Park Soil Data:** Swann Park soils were sampled for arsenic and kepone in 1976. Forty-five surface samples (1-2" depth) were collected by Allied Chemical Corporation (1976a). Twelve additional surface (0-3") and six subsurface (3-9") cores were collected and analyzed (Allied Chemical, 1976b), and four composite samples of the baseball and softball fields were also collected and analyzed (Allied Chemical, 1976c). The geometric mean of the initial 45 surface arsenic samples was 313 mg/kg with a maximum value of 6,600 mg/kg (located in the northwest corner of the Park). The spatially composited (averaged) arsenic samples of the four

ball fields had much lower concentrations (24 to 144 mg/kg) with the northwest field having the highest arsenic value.

The high concentrations of arsenic and kepone along the northern boundary of the Park led the ‘State Kepone Task Force’ to recommend that the 100 foot strip of park paralleling the Allied Chemical Corporation boundary be plowed, raked, top-soiled, and sodded (Allied Chemical, 1976d). The ‘State Kepone Task Force’ reviewed the soil sampling data as well as air sampling data and biological analyses of blood kepone concentrations and determined that ‘‘there are no significant health hazards existing in Swann Park...’’ Based on comments from Park personnel, ATSDR has determined that these recommendations were completed as specified.


Following a 2007 inquiry by Honeywell to MDE concerning the results of the 1976 ‘‘State Kepone Task Force,’’ Honeywell collected 25 surface and subsurface soil samples from Swann Park, and analyzed these samples for arsenic, hexavalent chromium, and kepone. A subset of those samples were further analyzed for a broad array of metals, pesticides, and semi-volatile organic compounds (SVOCs; Honeywell, 2007a/b). These are the soil data evaluated in this assessment. Following receipt of these results, Swann Park was closed pending clean up. Honeywell, with oversight from MDE and the City of Baltimore, is currently developing an extensive re-sampling program to continue the environmental evaluation of Swann Park.

The Swann Park Community: Figure 1 shows the location of Swann Park and the surrounding community. This figure also shows how many people live within one half mile of the site. Swann Park is located in an industrial area of south Baltimore and is bounded by the Interstate 95 and the Patapsco River. Based on the 2000 census, approximately 3,650 people live within one half mile of the park, with most of those homes located north of Interstate 95. Although relatively few people live directly adjacent to the park, it is highly used as a recreational area.

**Figure 1. Location of Swann Park and surrounding community**  
Baltimore, MD



EPA Facility ID: UNAVAILABLE



Site Location: Baltimore City County, MD





**Demographic Statistics**  
Within Area of Concern\*

Total Population	3,650
White Alone	3,285
Black Alone	236
Am. Indian & Alaska Native Alone	17
Asian Alone	42
Native Hawaiian & Other Pacific Islander Alone	0
Some Other Race Alone	23
Two or More Races	47
Hispanic or Latino**	69
Children Aged 6 and Younger	328
Adults Aged 65 and Older	401
Females Aged 15 to 44	826
Total Housing Units	1,622

Base Map Source: Geographic Data Technology, May 2005.  
Site Boundary Data Source: ATSDR Geospatial Research, Analysis, and Services Program, Current as of Generate Date (bottom left-hand corner).  
Coordinate System (All Panels): NAD 1983 StatePlane Maryland FIPS 1900 Feet

Demographics Statistics Source: 2000 U.S. Census  
\* Calculated using an area-proportion spatial analysis technique  
\*\* People who identify their origin as Hispanic or Latino may be of any race.



project=MDSAID62982-1;user=JACD-1;geo=Baltimore County, MD;keywords=Swan Park



FOR INTERNAL AND EXTERNAL RELEASE  
AGENCY FOR TOXIC SUBSTANCES AND DISEASE REGISTRY | UNITED STATES DEPARTMENT OF HEALTH AND HUMAN SERVICES

## Results

### Contaminants of Concern

ATSDR screened all of the contaminants detected in the April 2007 Swann Park soil samples against environmental guidelines. Of the many contaminants analyzed and detected in Swann Park soils (24 metals, 38 pesticides, and 65 SVOCs), only arsenic was consistently detected above its environmental guideline (20 mg/kg; Attachment 1).

The environmental guideline for lead is based on the average concentration in a residential yard or child's play area (400 mg/kg; EPA, 1998). The upper 95<sup>th</sup> percentile confidence limit of the geometric mean for lead is 271 mg/kg in Swann Park.<sup>1</sup> Most lead measurements were well below the 400 mg/kg screening value as was the average lead concentration (Attachment 1). Due to the limited distribution of lead concentrations above this screening level, ATSDR determined that widespread and repetitive exposures to the lead detected at the park are unlikely. Consequently, ATSDR screened out lead in Swann Park soil as a contaminant of concern at this site, and did not calculate site-specific lead exposure doses.

Arsenic was detected in most of the Swann Park samples (Honeywell, 2007b; Attachment 1). Concentrations in surface samples (0 to 3" depth) ranged from 37 to 1,330 mg/kg. The geometric mean of the 25 arsenic surface samples was 133 mg/kg with a maximum value of 1,330 mg/kg (Table 1; located in the northwest corner of the Park). Note that Table 1 also lists the arsenic and kepone concentrations from the 1976 samples.

Kepone (chlordecone) was detected in most of the recent soil samples from Swann Park. Although kepone concentrations in soil were well below the environmental guideline (30 mg/kg; Attachment 1), potential doses were estimated because it was processed at the Race Street facility and has widespread distribution in Swann Park soils. Kepone concentrations had a geometric mean of 0.2 mg/kg with a maximum value of 1.6 mg/kg (Table 1). Note that there are no available screening values for several other analytes listed in attachment 1. However, as these analytes were not specifically manufactured or processed at the Race Street facility, they are not likely to be present in significant concentrations in Swann Park soils and are not further evaluated in this health consultation.

Several poly aromatic hydrocarbons (PAHs; e.g. anthracene, benzo(a)anthracene, benzo(a)pyrene, and benzo(b)fluoranthene) were also detected. However, most detections were at very low concentrations (< 0.1 mg/kg) and the maximum concentrations were relatively low (2-5 mg/kg). Because the average concentrations of these PAHs are below their respective screening values and they were not specifically manufactured or processed at the Race Street

---

<sup>1</sup> There were five individual lead measurements above the environmental guideline (400 mg/kg) and only two of those were in the surface soils (three were in subsurface soil). Note that the 400 mg/kg environmental guidance level for lead in residential soils refers to an average concentration and discrete measurements are not directly comparable with this environmental guideline. Average soil concentrations are used for screening and dose assessment because exposure to soil occurs over a large area and duration of time. The exception is for pica children who may be exposed during a single event to a localized area of soil.

facility they will not be further evaluated in this health consultation. These analytes will be further evaluated if future samples indicate higher or more widespread distributions.

**Table 1. Summary of arsenic and kepone surface soil concentrations from Swann Park.**

Contaminant	Concentration Range (mg/kg)		Geometric Mean (mg/kg)		95 <sup>th</sup> UCL of Geomean	
	1976	2007	1976	2007	1976	2007
Arsenic	12—6,600	37—1,330	313	139	486	210
Kepone	1.5—25	0.05—1.6	8.1	0.2	10.2	0.3

### Exposure Scenarios

Calculation of exposure doses requires a specific estimate of the activities and time for each type of exposure. This evaluation will rely on the types of park activities described in the above *Community Health Concerns* to identify the types of exposures that occurred at Swann Park. Specifically, we evaluated exposures related to the following.

- Youth (teen) athletic activities (such as baseball, football, and other sports)
- Park workers
- Adult athletic coaches
- Adult park visitors and spectators (including nearby residents who may have visited the park on a regular basis)
- Children playing at the Park (including nearby residents who may have visited the park on a regular basis)
- Pica children (Geophagy, including pica behavior, refers to the intentional ingestion of soil items. The sensitive population associated with pica behavior is children aged 1–3 years old.)

Each of the above activities or exposure scenarios involves different behaviors and timeframes of exposure to soil. Table 2 lists the exposure scenarios and the assumed durations, soil intake rates, body weights, and time frame for each type of exposure. The primary exposure route of concern for surface soil is ingestion (oral exposure). This ingestion could occur by the inadvertent consumption of soil on hands or food items, mouthing of objects, or intentional ingestion (pica behavior).

The assumed exposure durations (Table 2) are based on discussions with community members, coaches, and park workers. Soil intake rates and body weights are based on recommendations

from the EPA Exposure Factor Handbook (EPA, 1999) and the ATSDR Public Health Assessment Guidance Manual (ATSDR, 2005a). Soil intake rates for teen athletes and park worker are based on intake rates for adult agricultural workers (EPA, 1999).

**Table 2. Exposure scenarios and assumptions for people using Swann Park.**

<b>Exposure Scenario</b>	<b>Duration of Exposure</b>	<b>Soil Intake Rate</b>	<b>Body Weight</b>	<b>Time Frame for Cancer Evaluation*</b>
Toddler (pica child)	A few hours to a day	5,000 mg/day 0.7 tsp/event	10 kg (22 lbs.)	NA; Single event; Acute Exposure
Child (non-pica)	365 days/yr	100 mg/day 0.02 tsp/day	14 kg (22 lbs.)	6 years
Adult Visitor or Spectator (including nearby residents)	365 days/yr	50 mg/day 0.01 tsp/day	70 kg (154 lbs.) Adult men/women body wt.	4 years
Teen Athlete	365 days/yr	100 mg/day 0.02 tsp/day	50 kg (110 lbs.) Average 13 yr old child	4 years
Adult Coach	365 days/yr	100 mg/day 0.02 tsp/day	70 kg (154 lbs.) Adult body wt.	20 years
Adult Park Worker	365 days/yr	100 mg/day 0.02 tsp/day	80 kg (176 lbs.) Adult male body wt.	30 years
Soil intake rates: Teen athlete, coach, and adult park worker soil ingestion rates are based on the adult agricultural worker soil intake rate (EPA Exposure Factors Handbook), others are ATSDR or EPA recommended rates (central tendency) for children and adults (EPA Exposure Factors Handbook). The soil intake rates, as converted to teaspoons, are based on a soil bulk density of 1.5 g/cm <sup>3</sup> and a volumetric conversion of 1 tsp = 4.93 cm <sup>3</sup> . * Estimates of excess lifetime cancer risk can be summed across scenarios if a person had exposures across multiple exposure scenarios (e.g., a person who played at the park as a young child, then played sports on the fields as a student athlete, and then visited the park later as an adult).				

### Estimated Exposure Doses

Estimating an exposure dose requires identifying how much, how often, and how long a person may come in contact with some concentration of the chemical in a specific medium (air, water, soil). The equation and assumptions used to estimate exposure doses from ingesting arsenic in surface soil follow (ATSDR, 2005a).

#### Equation 1: Exposure Dose Equation for Ingestion

$$\text{Dose} = \frac{\text{C} \times \text{IR} \times \text{EF} \times \text{CF} \times \text{BF}}{\text{BW}}$$

Where:

D = exposure dose in milligrams per kilogram per day (mg/kg/day)

C = chemical concentration in milligrams per kilogram (mg/kg)

IR	=	intake rate in milligrams per day (mg/day)
EF	=	exposure factor (unitless = 1)
CF	=	conversion factor, $1 \times 10^{-6}$ kilograms/milligram (kg/mg)
BF	=	bioavailability factor (unitless)
BW	=	body weight in kilograms (kg)

For arsenic in soil, the primary exposure route of concern is ingestion (oral exposure). Ingestion of soil could occur by the inadvertent consumption of soil on hands or food items, inhalation and subsequent ingestion of soil particles in the air, mouthing of objects, or intentional ingestion (pica behavior). For kepone exposures, the rate of absorption through skin (dermal absorption) must also be considered. The dermal dose equation is similar to the ingestion dose equation (1) except that soil adhered to skin is substituted for the intake rate (Table 3). Estimates of soil adhered are based on activity-specific soil adherence rates from the EPA Exposure Factors Handbook (EPA, 1999, Table 6-12). Note that these soil adherence rates are very similar to the soil ingestion rates (Table 2). Dermal doses are not important for arsenic and lead because these metals are not readily absorbed through the skin.

**Table 3. Soil adherence rates for kepone dermal dose estimation.**

<u>Scenario</u>	<u>Soil Adhered mg/day</u>
Adult Park Worker	427.5
Teen Athlete	461.7
Toddler (Pica Child)	356.3
Child (non pica)	356.3
Adult Visitor	100
Adult Coach	200

These soil adherence rates are derived by multiplying activity-specific rates of soil adherence ( $\text{mg}/\text{cm}^2$ ) with the recommended surface area ( $\text{cm}^2$ ) for outdoor soil contact (EPA Exposure Factors Handbook Table 6-12 and 6-14, respectively; EPA, 1999).

ATSDR estimated surface soil exposure doses using Equation 1. In the absence of complete exposure-specific information, ATSDR applied several health protective exposure assumptions to estimate exposures as accurately as possible. Specifically, ATSDR estimated arsenic exposure doses using the following assumptions and default intake rates for exposure via ingestion:

- The contaminant concentrations used in estimating the dose for the child exhibiting pica behavior are the maximum detected values (arsenic—1330 mg/kg, kepone—1.6 mg/kg). All other doses are the upper 95<sup>th</sup> percentile confidence limits of the geometric means (arsenic—210 mg/kg, kepone—0.30 mg/kg).<sup>1</sup> These 95% UCL values are health protective estimates of the average soil concentrations.

---

<sup>1</sup> The geometric mean is the mean or average of the logarithms of the concentrations. Log transformation of contaminant concentrations is applied because these geological parameters are log-normally distributed. This type of skewed distribution means that there are typically many low values with a few high values. The log-transformed concentrations are approximately normally distributed. The upper 95<sup>th</sup> percentile confidence limit means that there is a 95% chance that the average concentration is less than the upper confidence limit value.

- Soil intake rates, exposure factors, and body weights for all exposure scenarios are as listed in Tables 2 and 3.
- The bioavailability (BF) of arsenic was assumed to be 50% for all exposure scenarios except pica behavior (100%) due to direct effects on the gastrointestinal system. Most studies indicate that arsenic bioavailability is in the range of 15-40%. The bioavailability of arsenic for pica behavior and kepone was assumed to be 100%—that is, all of the contaminant ingested or adhered to skin was assumed to enter the bloodstream.
- Arsenic is excreted rapidly from the body such that for non-cancer effects there is no carryover from one sport season to the next. For cancer effects, the doses from multiple season and multiple year exposures were summed to assess long term exposures.

ATSDR derived exposure doses are based on these health protective assumptions. The estimated doses for arsenic and kepone exposures (based on 2007 soil data) are listed in Table 4. Estimated kepone doses for all exposure scenarios are at least 10 times lower than the respective health guidelines (Table 4). On the basis of these results, exposures to kepone at Swann Park are unlikely to result in any adverse health effects and will not be discussed in the following section on *Public Health Implications*.

Most of the calculated arsenic doses are also below their respective health guidelines (Table 4). However, arsenic exposures for pica children are greater than the short term (acute) minimal risk level (MRL). It is important to emphasize that a calculated dose greater than the MRL does not necessarily mean that a child consuming 5 grams (~1 teaspoon) of Swann Park soil will become sick. The potential health effects of a community member's exposure to arsenic are detailed in the following section.

Arsenic is a known human carcinogen and has been linked with skin, bladder, lung, and other cancers (ATSDR, 2005b). Consequently, ATSDR also evaluated the potential for Swann Park arsenic exposures to cause cancer by estimating a theoretical excess lifetime cancer risk level (ATSDR, 2005a).

### **Equation 2. Excess Lifetime Cancer Risk**

$$\text{Cancer risk} = \text{dose}(\text{mg/kg/day}) \times \text{cancer slope factor}(\text{mg/kg/day})^{-1}$$

These cancer risks are usually compared with a  $10^{-6}$  risk (one in a million) as defined by other governmental agencies. As with the non-cancer evaluation approach, exposures below a screening value (by convention for cancer risk evaluations, less than a  $10^{-6}$  risk) require no further evaluation, while estimated dose levels that exceed the  $10^{-6}$  value are evaluated further. The potential for observing adverse effects is made on the basis of dose evaluation (a margin of exposure or MOE approach), rather than on the basis of theoretical risk calculations.

**Table 4. Calculated doses for arsenic and kepone at Swann Park; 2007 soil data.**

Exposure Scenario	Contaminant	Estimated Dose (mg/kg/day)	Health Guideline (mg/kg/day) Source	Theoretical Excess Lifetime Cancer Risk
Toddler (pica child)	Arsenic	3.3E-01	5.0E-03 Acute MRL	Not Applicable
	Kepone	8.6E-04	1.0E-02 Acute MRL	
Child (non-pica)	Arsenic	7.5E-04	3.0E-04 Chronic MRL	9.7E-05
	Kepone	1.4E-05	5.0E-04 Chronic MRL	7.5E-06
Teen Athlete	Arsenic	2.1E-04	3.0E-04 Chronic MRL	1.8E-05
	Kepone	3.4E-06	5.0E-04 Chronic MRL	1.6E-06
Adult Coach	Arsenic	1.5E-04	3.0E-04 Chronic MRL	6.4E-05
	Kepone	1.3E-06	5.0E-04 Chronic MRL	4.0E-06
Adult Visitor	Arsenic	7.5E-05	3.0E-04 Chronic MRL	6.4E-06
	Kepone	6.5E-07	5.0E-04 Chronic MRL	2.0E-07
Adult Park Worker	Arsenic	1.3E-04	3.0E-04 Chronic MRL	8.4E-05
	Kepone	2.0E-06	5.0E-04 Chronic MRL	6.9E-06

- Doses are calculated using the 95% upper confidence limit of the geometric mean of the contaminant concentrations and the exposure scenarios from Table 2 (pica uses maximum).
- Because kepone may be absorbed through skin contact, kepone doses include an ingestion component plus a dermal absorption component.
- Excess lifetime cancer risk is not applicable to short term pica behavior.
- Excess lifetime cancer risks for multiple exposure scenarios can be summed if a person used the park for multiple activities. For example, if someone visited the park as a child and a teen athlete, the arsenic cancer risk would be  $9.7E-05 + 3.6E-05 = 13.3E-05$  (or  $1.33E-04$ ).
- MRL—Minimal risk level; An ATSDR estimate of daily human exposure to a hazardous substance at or below which that substance is unlikely to pose an appreciable risk of harmful (adverse), non-cancerous effects. MRLs are calculated for a route of exposure (inhalation or oral) over a specified time period (acute, intermediate, or chronic).
- These MRLs are all for oral or ingestion exposure; also the chronic and intermediate MRLs for kepone have the same value.

Arsenic and kepone doses estimated from the 1976 soil data would be higher than exposure doses estimated from the 2007 data since the reported soil concentrations were higher. Also, during the time the Race Street plant was operating, there may have been an additional inhalation dose from airborne emissions from the plant. Consequently, we cannot determine the potential hazard of arsenic and kepone doses at Swann Park when the Race Street Plant was operating (prior to 1976).

## Public Health Implications

Arsenic occurs naturally in soil and minerals. People normally take in small amounts of arsenic in air, water, soil, and food. Of these, food is usually the most common source of arsenic for people (ATSDR 2005). In order to determine whether the potential exposures to arsenic-contaminated soil presents a public health hazard at this site, ATSDR compared the estimated doses with benchmarks or screening doses that are derived from dose levels known to produce adverse health effects. For arsenic, ATSDR has developed minimal risk levels (MRLs) that cover brief exposures (acute, or less than 14 days) and longer term exposures (chronic, or more than a year).

An MRL is an estimate of the daily human exposure to a hazardous substance that is likely to be without appreciable risk of adverse non-cancer health effects over a specified duration of exposure. The MRL is derived from exposure levels observed to produce adverse effects, with uncertainties (or safety factors) incorporated into the value. Thus, MRLs are intended only to serve as a screening tool to help public health professionals decide which exposure situations require more extensive evaluation. Estimated exposure dose levels below an MRL are not likely to produce non-cancer adverse effects. Exposure estimates above an MRL do not mean that adverse effects will occur, but rather that further evaluation of the exposure is warranted. ATSDR then evaluates the potential for adverse health effects in an exposed community by comparing levels known to produce adverse effects to the estimated site-related doses. This margin of exposure (MOE) approach, along with an evaluation of available epidemiologic, toxicologic, and medical data, is used by health assessors as part of the public health determination to reach qualitative (rather than quantitative) decisions about hazards posed by site-specific conditions of exposure.

At low-level exposures, arsenic compounds are detoxified—that is, changed into less harmful forms—and excreted in the urine (ATSDR, 2005). At higher-level exposures, however, the body may not have the ability to detoxify the increased amount of arsenic. When this overload happens, blood levels of arsenic increase and adverse health effects may occur. Arsenic, like some other chemicals, does not seem to cause adverse health effects until a certain amount, or threshold, of the chemical has entered the body. Once the threshold, also known as the minimal effective dose, is reached adverse health effects may result (ATSDR 2005b).

ATSDR reviewed the scientific literature regarding arsenic toxicity to evaluate whether non-cancer adverse health effects would be expected to occur at the estimated exposure doses. The acute oral MRL for arsenic (0.005 mg/kg/day) is based on several temporary effects that could occur from acute exposures ( $\leq 14$  days). Acute exposure to arsenic can be toxic to the stomach and intestines, with symptoms such as pain, nausea, vomiting, and diarrhea. When an estimated acute exposure dose for pica behavior is below 0.005 mg arsenic/kg/day, non-cancerous effects are unlikely. It should be noted that the acute MRL is 10 times below the levels reported to cause these effects in humans (acute Lowest Observed Adverse Effect Level (LOAEL) = 0.05 mg/kg/day).

The estimated exposure dose for children exhibiting pica behavior at Swann Park exceeds the ATSDR acute MRL, and is also above the LOAEL. Sensitive populations, such as children who eat non-food items like soil, could receive doses that might cause temporary effects (e.g., nausea, diarrhea, vomiting and stomach cramps). Groups that are at an increased risk for pica behavior are children aged 1–3 years old. ATSDR suggests that concerned parents monitor their children's behavior while they are playing outdoors to ensure that their children are not exhibiting pica behavior and eating excessive amounts of soil and discuss their concerns and/or observed behaviors with their doctor.

In addition to the acute MRL, ATSDR developed a chronic oral MRL for arsenic of 0.0003 mg/kg/day. The estimated exposure doses for a child (non-pica) are above the chronic oral MRL (see Table 4). The estimated exposure doses for teen athlete, adult visitor, adult coach, and adult worker do not exceed the chronic oral MRL. The ATSDR chronic oral MRL is based on common and characteristic effects of arsenic ingestion; a pattern of skin changes that include hyperpigmentation and hyperkeratosis. These dermal effects have been noted in some human studies that involved daily, long-term ingestion (more than 45 years) of elevated arsenic levels in drinking water. Collectively, these studies indicate that the dose for hyperpigmentation and hyperkeratosis is 0.014 mg As/kg/day (ATSDR 2005b). No adverse health effects have been observed (NOAEL) at arsenic doses of 0.008 mg/kg/day.

The estimated chronic (long term) doses for a child (non-pica) exposure scenario is below the arsenic NOAEL dose of 0.008 mg/kg/day. Additionally, the exposure doses were calculated assuming continuous daily exposure (365 days/yr for 6 years). The calculated doses are a health protective overestimate as children were not exposed everyday over several years. Studies have shown that 45 to 85 percent of an ingested dose of arsenic is eliminated within 1 to 3 days; however, some remains for several months or longer (Buchet et al. 1981; Crecelius 1977; Mappes 1977; Tam et al. 1979). Therefore, we do not expect that chronic exposure to arsenic in Swann Park soil would result in a dose level or body burden high enough to cause adverse health effects. Based on a review of this information, ATSDR concludes that chronic non-cancer health effects are not expected to occur in children or adults based on the 2007 soil sampling results.

Regarding historical chronic exposure to arsenic and kepone in Swann Park soil, we used the upper 95<sup>th</sup> percentile confidence limits of the geometric means (arsenic—486.4 mg/kg, kepone—10.2 mg/kg) from the 1976 reported soil concentrations (see Table 1) under the same exposure scenarios and default assumptions used above. The estimated chronic doses from kepone exposure in soil for the teen athlete and adult coach scenarios were just above the health guideline value and the park worker dose was equal to the health guideline (Table 4). Although these estimated doses for chronic exposure to arsenic in soil are above the screening value, they are more than an order of magnitude below levels shown to cause dermal effects. Therefore, chronic non-cancer health effects are not expected to occur for any exposure scenarios based on the 1976 soil sampling results.

The estimated exposure dose (1976 data) for children exhibiting pica behavior at Swann Park exceeds the ATSDR acute arsenic and kepone MRLs, and is also above the LOAEL for arsenic. Sensitive populations, such as children who eat non-food items like soil, could receive arsenic doses that might cause temporary effects (e.g., nausea, diarrhea, vomiting and stomach cramps).

For kepone, the estimated pica child exposure dose (0.00016 mg/kg/d) was well below the acute MRL (0.01 mg/kg/day).

ATSDR discussed concerns about this site with Dr. Genevieve Matanoski, a researcher from Johns Hopkins University. Dr. Matanoski conducted an investigation of environmental data and cancer information for the site area (Matanoski et al., 1981). She found the distribution of arsenic in soil near the old Allied plant and along the railroad line was higher than in control areas in the city, and based on a review of death certificates concluded that men living in close proximity to the old Allied facility had a higher risk of lung cancer in the 1960s and early 1970s compared to individuals in other areas of the city. Dr. Matanoski emphasized to ATSDR that the key issue regarding current exposures at Swann Park is the bioavailability of the arsenic in the soil to users of the playing fields. Consistent with the conclusions in this document, Dr. Matanoski stated that it would be very difficult for users of the park to swallow enough dirt from activities at the park to achieve a dose with toxic health effects. She stated that she believes that her original findings about elevated cancer risk were related to ongoing air exposures from the Allied Chemical facility, and not from incidental soil exposures.

Excess lifetime cancer risks for exposure to Swann Park soils are presented in Table 4. These theoretical risks assume that a park worker has been exposed to Swann Park soils for 30 years, an adult coach has been exposed to Swann Park soils for 20 years, and community members were exposed to Swann Park soils as young children who played in the park as a child for six years, then played sports in the park as a student athlete for 4 years, and visited the park as an adult for 4 years. The resulting excess lifetime cancer risks from exposure to arsenic in Swann Park soils are all less than 1 in 10,000 (9.7E-05; Table 4). Note that these excess lifetime cancer risks can be summed if a person used the park for multiple activities.

The excess cancer risks (Table 4) represent the theoretical increase in cancer risk due to exposure to arsenic in Swann Park soils. All of the uncertainties and health-protective assumptions associated with the dose calculations are included in the risk estimations, as well as the uncertainty in deriving the cancer slope factor (EPA, 2003). These estimates of cancer risk can be considered suggestive of a *low to no apparent increased risk*. These low risk estimates combined with the likely overestimation of the arsenic doses indicate that exposure to Swann Park soil is not likely to result in excess cancer in people who have used Swann Park.

Note that the above estimates of excess lifetime cancer risk are based on dose estimates from the 2007 soil sampling data. In the absence of air monitoring data from plant emissions, we cannot speculate on potential cumulative arsenic and kepone doses and cancer risks at Swann Park when the plant was operating (prior to 1976).

The arsenic doses and derived cancer risks presented in Table 4 are based on measured soil concentrations sampled in 2007. The sampling procedures used in both sampling events (1976 and 2007) have significant limitations in determining whether isolated areas (hot spots) of high contaminant concentrations are present. If present, the spatial extent and average concentrations of such hot spots are unknown. Based on the 1976 sampling (Allied Chemical, 1976c), it does appear that the northwest softball field had a higher average arsenic concentration than the other fields. However, the composited average concentration of this field (144 mg/kg) was much

lower than the 95<sup>th</sup> UCL (210 mg/kg) used in our dose calculations. These composited 1976 arsenic concentrations suggest that we have greatly overestimated potential arsenic doses. Ongoing soil sampling should reduce the uncertainty in these dose calculations.

### **Child Health Concerns**

In communities faced with air, water, or food contamination, the many physical differences between children and adults demand special emphasis. Children could be at greater risk than are adults from certain kinds of exposure to hazardous substances. Children play outdoors and sometimes engage in hand-to-mouth behaviors that increase their exposure potential. Children are shorter than are adults; this means they breathe dust, soil, and vapors close to the ground. A child's lower body weight and higher intake rate results in a greater dose of hazardous substance per unit of body weight. If toxic exposure levels are high enough during critical growth stages, the developing body systems of children can sustain permanent damage. Finally, children are dependent on adults for access to housing, for access to medical care, and for risk identification. Thus adults need as much information as possible to make informed decisions regarding their children's health.

There is a significant amount of toxicological data available on arsenic and human health effects. However, the data needed to account for an accurate representation of any potential unique concerns related to early-life exposures to arsenic appears to be insufficient. The National Research Council concluded in 2001 that "There are no reliable data that indicate heightened susceptibility of children to arsenic" (NRC, 2001).

The special concerns related to a child's health have been addressed in this assessment by specifically using exposure scenarios based on a child's soil intake, body weight, and park activities. The previous discussion of potential health effects from arsenic exposure also explicitly includes health endpoints in children. Children that may have consumed teaspoon quantities of Swann Park soil may have experienced temporary symptoms such as nausea, diarrhea, vomiting and stomach cramps due to arsenic exposure. These short term exposures are unlikely to result in any other long term symptoms or diseases.

### **Response to Community Concerns**

This assessment of exposures to arsenic-contaminated soil at Swann Park has attempted to incorporate the communities health concerns into the specific health issues that have been addressed as well as the types of exposures that have been evaluated. In order to highlight how ATSDR has responded to those concerns, Table 5 lists the specific concerns collected during public meetings along side the ATSDR response to each concern. ATSDR welcomes additional comments or concerns and will further address any unresolved issues as appropriate.

**Table 5. Community concerns related to Swann Park and ATSDR responses.**

Community Concern	ATSDR Response
<p>Inhalation of dust that would blow from the park towards their homes</p>	<p>Inhalation of dust is a potential route of exposure to contaminated soil. However, even though the dust is initially inhaled, most soil particles larger than ~1 micron, which includes essentially all re-suspended soil particles, end up in the digestive tract and are evaluated as part of the ingested soil. Although a small fraction of the arsenic in inhaled dust may be absorbed in lungs, it would be a minor fraction compared to the ingested arsenic. Current arsenic concentrations in nearby homes/yards are expected to be much lower than levels in the park and are unlikely to present a public health hazard. In the absence of historic air monitoring data, ATSDR cannot speculate on airborne arsenic concentrations when the Race Street plant was operating (prior to 1976).</p>
<p>Relationship between exposure to arsenic-contaminated soils and skin conditions, seizures in a toddler, nervous system/mental health conditions, sleep apnea, diabetes, and a fatal blood clot/aneurysm</p>	<p>Arsenic exposure has been related to several skin conditions including hyperpigmentation, hyperkeratosis, and skin cancer. The doses at which those diseases occur are much higher than any of the doses likely experienced at Swann Park. Also acute, high-dose exposures (much higher than those estimated for Swann Park) often lead to encephalopathy, with signs and symptoms such as headache, lethargy, mental confusion, hallucination, seizures, and coma (ATSDR 2005). An association has been demonstrated between exposure to arsenic in drinking water and increased incidence of diabetes mellitus (ATSDR 2005), although dose response relationships are not available and the mechanism of action for this response has not been characterized. There is little evidence to suggest that arsenic functions as an endocrine disruptor. ATSDR is not aware of any studies or associations between arsenic-contaminated soils and any of the other health conditions mentioned (seizures, sleep apnea, or blood clots).</p>
<p>Medical testing options for individuals who were exposed to the contamination at the park and guidance to parents who have already pursued getting hair and nail medical testing for their children.</p>	<p>Based on the results of this assessment, exposure to arsenic contaminated soil at Swann Park does not present a short or a long term public health hazard likely to produce health effects in individual users of the park, with the exception of young toddlers visiting the park exhibiting pica behavior. Consequently, medical testing of individuals is not recommended and any arsenic-specific test results may have little relation to Swann Park exposures. Community members are referred to Attachment 1, which includes information about the limitations of hair testing generally. Community members who have already pursued hair testing are welcome to consult directly with ATSDR or have their physicians consult with ATSDR regarding interpretation of their results.</p>
<p>Long term follow up for people who were exposed to contamination at the park</p>	<p>As above, adverse health effects from Swann Park arsenic exposures are unlikely and long term medical follow-up is unnecessary.</p>

**Table 5. Community concerns related to Swann Park and ATSDR responses.**

<b>Community Concern</b>	<b>ATSDR Response</b>
Soil exposures to student athletes and their coaches (including field maintenance activities, lack of water, hand washing facilities, and baseball catchers, etc.)	This evaluation explicitly included exposure assumptions related to student athletes and coaches. These assumptions include high soil ingestion rates based adult agricultural worker activities (and dermal contact rates for soccer players and groundskeepers for kepone dermal exposures). Arsenic is excreted rapidly from the body such that for non-cancer effects there is no carryover from one sport season to the next. For cancer effects, the doses from multiple season and multiple year exposures are assumed to occur 365 days per year for four years.
What are the health effects from ingestion of arsenic-contaminated soil? How much soil would produce toxic effects?	The potential health effects from exposure to arsenic are presented in the preceding section. Table 2 presents ingestion rates in terms of teaspoons of soil (assuming a soil density of 1.5 g/cm <sup>3</sup> ).

## **Conclusions, Recommendations, and Public Health Action Plan**

### **Conclusions**

Recent analyses of surface soil at Swann Park have measured concentrations of arsenic that exceed various environmental guidelines. Other analytes, such as chromium, PAHs, and kepone were also detected but at levels that do not exceed such guidelines. Normal park activities, such as field sports (baseball, football, etc.), and park maintenance have resulted in exposures to the arsenic-contaminated soil. With the exception of pica children, estimates of arsenic exposure doses based on these specific park activities result in doses that are not likely to produce adverse health effects. A child consuming about a teaspoon of soil may experience temporary nausea, diarrhea, vomiting or stomach cramps. These acute effects are unlikely to continue and other long term adverse health effects, including cancer, are unlikely for all exposures.

Dose estimates based on historic soil sampling present the same conclusions as the recent soil data. That is, with the exception of pica children, estimates of arsenic doses based on the above park activities are not likely to produce adverse health effects. However, there may have been an additional dose component from inhalation of airborne plant emissions when the plant was operating. Consequently, the conclusions of this health consultation are limited to the time period.

### **Recommendations**

1. Communicate the findings from this Health Consultation to the concerned community. This should include a presentation at a public meeting with the City of Baltimore Health Department, an individualized meetings with nearby residents, and outreach directly to Digital Harbor High School (e.g., to the school nursing staff and/or athletic department).
2. Concerned community members are encouraged to share these findings with their personal physicians. Community members who have already pursued hair testing are welcome to consult directly with ATSDR or have their physicians consult with ATSDR regarding interpretation of their results.
3. Additional sampling should be conducted to ensure that the data reviewed in this consultation are representative of the entire park and that there are no isolated hot spots of high arsenic concentrations. ATSDR encourages Honeywell and/or MDE to consider evaluation of the potential for migration of contamination to the block of residences

**Public Health Action Plan** located near the park in pending site characterization activities.

1. ATSDR will present the results of this health consultation to the Swann Park community and respond to ongoing public health concerns upon completion.
2. Honeywell, with guidance from MDE and the City of Baltimore, will develop and conduct an extensive re-sampling effort of Swann Park.
3. ATSDR will provide comments on the proposed sampling plan and evaluate the resulting data and revise the conclusions of this consultation accordingly.

### **Authors**

Mark W. Evans, Ph.D.  
Geologist, Site Radiological and Assessment Branch  
Division of Health Assessment and Consultation  
Agency for Toxic Substances and Disease Registry

Karl Markiewicz, Ph.D.  
Toxicologist, ATSDR Region 3  
Division of Regional Operations  
Agency for Toxic Substances and Disease Registry

Lora Siegmann Werner, MPH  
Senior Regional Representative, ATSDR Region 3  
Division of Regional Operations  
Agency for Toxic Substances and Disease Registry

### **Reviewers**

Burt J. Cooper, MS  
Supervisory Environmental Health Scientist  
Site Radiological and Assessment Branch  
Division of Health Assessment and Consultation  
Agency for Toxic Substances and Disease Registry

Sandra G. Isaacs, Chief  
Site Radiological and Assessment Branch  
Division of Health Assessment and Consultation  
Agency for Toxic Substances and Disease Registry

Kenneth G. Orloff, Ph.D.  
Assistant Director for Science  
Division of Health Assessment and Consultation  
Agency for Toxic Substances and Disease Registry

## References

Allied Chemical, 1976a. Swann Park Soil Sampling. Baltimore Plant, Agricultural Division, Allied Chemical Corporation, February 25, 1976, provided by Honeywell, Baltimore MD.

Allied Chemical, 1976b. Swann Park-Core Samples. Baltimore Plant, Agricultural Division, Allied Chemical Corporation, March 18, 1976, provided by Honeywell, Baltimore MD.

Allied Chemical, 1976c. Swann Park Baseball Fields Soil Sampling. Baltimore Plant, Agricultural Division, Allied Chemical Corporation, March 19 and 23, 1976, provided by Honeywell, Baltimore MD.

Allied Chemical, 1976d. Allied Chemical Corporation Memorandum, State Kepone Task Force Meeting- March 24/April 19, 1976. Baltimore Plant, Agricultural Division, Allied Chemical Corporation, provided by Honeywell, Baltimore MD.

ATSDR, 2005a. Public Health Assessment Guidance Manual (Update). US Department of Health and Human Services, Agency for Toxic Substances and Disease Registry, Atlanta, GA, January 2005.

ATSDR, 2005b. Toxicological Profile for Arsenic, Draft for Public Comment. US Department of Health and Human Services, Agency for Toxic Substances and Disease Registry, Atlanta, GA, September 2005. <http://www.atsdr.cdc.gov/toxprofiles/tp2.html>

Baltimore Sun, 2007. "Arsenic forces closing of park; Tests show high levels in soil; city locks gates of Swann Park." 4/20/07 edition, Baltimore MD, accessed May 17, 2007. [http://www.baltimoresun.com/news/local/baltimore\\_city/bal-te.md.ci.arsenic20apr20,0,5283439.story?coll=bal-local-headlines](http://www.baltimoresun.com/news/local/baltimore_city/bal-te.md.ci.arsenic20apr20,0,5283439.story?coll=bal-local-headlines)

Buchet JP, Lauwerys R, Roels H. 1981. Comparison of the urinary excretion of arsenic metabolites after a single oral dose of sodium arsenite, monomethylarsonate or dimethylarsinate in man. *Int Arch Occup Environ Health* 48:71-9. Cited in: Agency for Toxic Substances and Disease Registry. 2000. Toxicological profile for arsenic. Atlanta: U.S. Department of Health and Human Services; September.

Crecelius EA. 1977. Changes in the chemical speciation of arsenic following ingestion by man. *Environ Health Perspect* 19:147-50. Cited in: Agency for Toxic Substances and Disease Registry. 2000. Toxicological profile for arsenic. Atlanta: U.S. Department of Health and Human Services; September.

EPA, 2003. Draft Final Cancer Risk Assessment Guidelines. [www.epa.gov/ncea/raf/cancer2003.htm](http://www.epa.gov/ncea/raf/cancer2003.htm) Washington, D.C., Environmental Protection Agency, February 28, 2003.

EPA, 1999. Exposure factors handbook. Washington, DC: Environmental Protection Agency, EPA/600/C-99/001; 1999 Feb. <http://www.epa.gov/ncea/pdfs/efh/front.pdf>

EPA, 1998. Lead; identification of dangerous levels of lead; notice of proposed rulemaking. U.S. Environmental Protection Agency. Fed Regist 63:30302.

Honeywell, 2007a. Draft Workplan for Soil Screening Assessment, Swann Park, Baltimore, Maryland. Prepared by CH2MHill for Honeywell, Morristown NJ, April 2007.

[www.mde.state.md.us/assets/document/SP\\_4-13-07\\_MDE\\_SwannPark\\_Soil\\_Sample\\_Screening\\_Work%20plan.pdf](http://www.mde.state.md.us/assets/document/SP_4-13-07_MDE_SwannPark_Soil_Sample_Screening_Work%20plan.pdf)

Honeywell, 2007b. Swann Park Results\_April2007.xls. EXCEL spreadsheet of soil sampling results prepared by CH2MHill for Honeywell, Morristown NJ, April 2007. Forwarded to ATSDR via e-mail (5/3/2007).

Mappes R. 1977. Experiments on excretion of arsenic in urine. Int Arch Occup Environ Health 40:267–72. Cited in: Agency for Toxic Substances and Disease Registry. 2000. Toxicological profile for arsenic. Atlanta: U.S. Department of Health and Human Services; September.

Matanoski et al., 1981. Cancer Mortality in an Industrial Area of Baltimore. Environmental Research 25, pp. 8-28 (1981).

MDE, 2007. Swann Park/Former Allied Chemical Race Street Site. Accessed May 2007. <http://www.mde.state.md.us/CitizensInfoCenter/Health/swannpark.asp>

National Research Council.(NRC). 2001. Arsenic in Drinking Water. 2001 Update. Washington, DC: National Academy Press.

Tam GKH, Charbonneau SM, Bryce F, Pomroy C, Sandi E. 1979. Metabolism of inorganic arsenic (74As) in humans following oral ingestion. Toxicol Appl Pharmacol 50:319–22. Cited in: Agency for Toxic Substances and Disease Registry. 2000. Toxicological profile for arsenic. Atlanta: U.S. Department of Health and Human Services; September.

**Attachment 1. Analytes and environmental guidelines for soil from Swann Park (from Honeywell, 2007).**

<i>Contaminant</i>	<i>Maximum mg/kg (ppm)</i>	<i>Geometric Mean mg/kg</i>	<i>Comparison Value mg/kg Source</i>
<b><u>Metals</u></b>			
Antimony	67	4.6	20 RMEG C
<b>Arsenic</b>	<b>1330 surface 2220 subsurface</b>	<b>156.1</b>	<b>20 EMEG CC</b>
Barium	167	84.7	30,000 EMEG CC
Beryllium	1.5	0.7	100 EMEG CC
Cadmium	1.3	0.3	10 EMEG CC
Chromium, Hexavalent	10.1	1.2	200 RMEG C
Cobalt	16.2	7.8	500 EMEG CI
Copper	116	46.4	500 EMEG CI
Lead	1080	196.7	400 EPA AL
Magnesium	8190	2322.5	NA
Manganese	537	214.1	3,000 RMEG C
Mercury	1.4	0.4	NA
Nickel	36.7	18.7	1,000 RMEG C
Potassium	2550	1253.9	NA
Selenium	2.2	1.3	300 EMEG CC
Silver	0.7	0.5	300 RMEG C
Thallium	2.1	1.7	NA
Vanadium	78.1	45.8	200 EMEG IC
Zinc	261	109.0	20,000 EMEG CC
<b><u>SVOCs - Method 8270C</u></b>			
1,1'-Biphenyl	0.05	0.04	3,000 RMEG C
2,2'-oxybis(1-Chloropropane)	0.05	0.04	NA
2,4,5-Trichlorophenol	0.10	0.08	5,000 RMEG C
2,4,6-Trichlorophenol	0.05	0.04	60 CREG
2,4-Dichlorophenol	0.05	0.04	NA
2,4-Dimethylphenol	0.10	0.08	1,000 RMEG C
2,4-Dinitrophenol	1.00	0.85	100 RMEG C
2,4-Dinitrotoluene	0.10	0.08	100
2,6-Dinitrotoluene	0.05	0.04	200 EMEG CI
2-Chloronaphthalene	0.05	0.04	4,000 RMEG C
2-Chlorophenol	0.05	0.04	300 RMEG C
2-Methylnaphthalene	0.14	0.05	200 RMEG C
2-Methylphenol	0.10	0.08	NA
2-Nitroaniline	0.05	0.04	NA
2-Nitrophenol	0.05	0.04	NA
3,3'-Dichlorobenzidine	0.15	0.13	NA
3-Nitroaniline	0.10	0.08	NA
4,6-Dinitro-2-methylphenol	0.26	0.21	NA
4-Bromophenyl-phenylether	0.05	0.04	NA

<i>Contaminant</i>	<i>Maximum mg/kg (ppm)</i>	<i>Geometric Mean mg/kg</i>	<i>Comparison Value mg/kg Source</i>
4-Chloro-3-methylphenol	0.10	0.08	NA
4-Chloroaniline	0.10	0.08	200 RMEG C
4-Chlorophenyl- phenylether	0.05	0.04	NA
4-Methylphenol	0.10	0.08	NA
4-Nitroaniline	0.10	0.08	NA
4-Nitrophenol	0.26	0.21	NA
Acenaphthene	0.51	0.05	3,000 RMEG C
Acenaphthylene	0.14	0.05	NA
Acetophenone	0.10	0.08	NA
Anthracene	1.20	0.07	20,000 RMEG C
Atrazine	0.05	0.04	200 EMEG CI
Benzaldehyde	0.10	0.08	5,000 RMEG C
Benzo(a)anthracene	2.90	0.29	NA
Benzo(a)pyrene	3.90	0.30	0.1 CREG
Benzo(b)fluoranthene	5.80	0.45	NA
Benzo(g,h,i)perylene	3.60	0.25	NA
Benzo(k)fluoranthene	1.70	0.16	NA
bis(2- Chloroethoxy)methane	0.05	0.04	NA
bis(2-Chloroethyl)ether	0.05	0.04	0.6 CREG
bis(2-Ethylhexyl)phthalate	0.36	0.14	NA
Butylbenzylphthalate	0.11	0.09	10,000 RMEG C
Caprolactam	0.05	0.04	30,000 RMEG C
Carbazole	0.56	0.05	NA
Chrysene	3.40	0.31	NA
Dibenz(a,h)anthracene	0.78	0.07	NA
Dibenzofuran	0.40	0.05	NA
Diethylphthalate	0.10	0.08	40,000 RMEG C
Dimethylphthalate	0.10	0.08	NA
Di-n-butylphthalate	0.10	0.08	5,000 RMEG C
Di-n-octylphthalate	0.10	0.09	20,000 EMEG CI
Fluoranthene	3.70	0.47	2,000 RMEG C
Fluorene	0.52	0.05	2,000 RMEG C
Hexachlorobenzene	0.05	0.04	5 EMEG CI
Hexachlorobutadiene	0.10	0.08	10 EMEG CI
Hexachlorocyclopentadiene	0.26	0.21	300 RMEG C
Hexachloroethane	0.05	0.04	50 RMEG C
Indeno(1,2,3-cd)pyrene	3.10	0.21	NA
Isophorone	0.05	0.04	10,000 RMEG C

<i>Contaminant</i>	<i>Maximum mg/kg (ppm)</i>	<i>Geometric Mean mg/kg</i>	<i>Comparison Value mg/kg Source</i>
Naphthalene	0.11	0.05	1,000 RMEG C
Nitrobenzene	0.05	0.04	30 RMEG C
N-Nitroso-di-n-propylamine	0.05	0.04	0.1 CREG
N-Nitrosodiphenylamine	0.05	0.04	0.01 CREG
Pentachlorophenol	0.26	0.21	50 EMEG CI
Phenanthrene	3.80	0.25	NA
Phenol	0.05	0.04	20,000 RMEG C
Pyrene	4.00	0.51	2,000 RMEG C
<b><u>Pesticides - Method 8081A</u></b>			
Kepone (chlordecone)	1.6	0.2	30 EMEG CC
Aldrin	0.01	0.001	2RMEG C
Alpha BHC	0.01	0.002	NA
Alpha Chlordane	0.01	0.002	30 RMEG C
Beta BHC	0.03	0.005	NA
Delta BHC	0.01	0.001	NA
Dieldrin	0.17	0.008	3 RMEG C
Endosulfan I	0.01	0.002	300 RMEG C
Endosulfan II	0.01	0.002	
Endosulfan Sulfate	0.01	0.002	
Endrin	0.01	0.002	20 RMEG C
Endrin Aldehyde	0.04	0.003	NA
Endrin Ketone	0.02	0.003	NA
Gamma BHC - Lindane	0.01	0.001	NA
Gamma Chlordane	0.03	0.007	30 RMEG C
Heptachlor	0.01	0.001	30 RMEG C
Heptachlor Epoxide	0.03	0.002	0.7 RMEG C
Methoxychlor	0.05	0.013	NA
p,p-DDD	0.37	0.025	3 CREG
p,p-DDE	8.40	0.447	2 CREG
p,p-DDT	2.90	0.218	30 RMEG C
Toxaphene	0.34	0.079	50 EMEG CI
<b><u>Pesticides 8141A &amp; Herbicides 8151A</u></b>			
2,4,5-T	0.03	0.001	500 RMEG C
2,4,5-TP	0.03	0.001	400 RMEG C
2,4-D	0.05	0.008	500 RMEG C
2,4-DB	0.01	0.007	NA
2,4-DP (Dichloroprop)	0.01	0.006	NA
Dalapon	0.07	0.023	NA
Diazinon	0.03	0.019	100 EMEG CI

<i>Contaminant</i>	<i>Maximum mg/kg (ppm)</i>	<i>Geometric Mean mg/kg</i>	<i>Comparison Value mg/kg Source</i>
Dicamba	0.03	0.003	2,000 RMEG C
Dinoseb	0.02	0.002	50 RMEG C
Ethion	1.00	0.042	30 RMEG C
Ethyl Parathion	2.70	0.048	NA
Guthion (Azinphos-methyl)	0.03	0.023	200 EMEG CC
Malathion	0.03	0.024	1,000 RMEG C
MCPA	4.90	1.555	NA
MCPP (Mecoprop)	1.20	0.630	NA
Methyl Parathion	0.03	0.028	10 RMEG C


Notes:

- Maximum and mean values for all analytes except arsenic and kepone include surface and subsurface samples.
- Contaminant maxima and geometric means include numeric values for contaminants with laboratory qualifiers of “U”. These contaminant values are considered as non-detections.
- NA: Comparison values (environmental guidelines) are not available for many contaminants.
- CREG: *Cancer risk evaluation guide* is the estimated concentration of a contaminant that would likely cause, at most, one excess cancer in a million people exposed over a lifetime. CREGs are calculated from cancer slope factors and health protective exposure assumptions.
- EMEG: *Environmental media evaluation guides* are contaminant concentrations calculated from “minimal risk levels” that are likely to be without an appreciable risk of adverse, non-cancer health effects for a specified duration of exposure. EMEGs are determined separately for children and adults and for various exposure durations.
  - EMEG CI refers to *child* exposures for an *intermediate* duration (14 to 365 days).
  - EMEG CC refers to *child* exposures for an *chronic* duration (more than 1 year).
- EPA AL: Environmental Protection Agency Action Limit (EPA, 1998).
- RMEG: *Reference dose media evaluation guide* are contaminant concentrations calculated from EPA Reference Doses that are likely to be without appreciable risk of adverse non-cancer health effects during a lifetime of exposure.
  - RMEG C refers to RMEGs calculated based on exposure assumptions for a *child*.

**Attachment 2. ATSDR Fact Sheet on Hair Sample Testing**

[http://www.atsdr.cdc.gov/HAC/hair\\_analysis/03-0330HairSampleTesting-Community.pdf](http://www.atsdr.cdc.gov/HAC/hair_analysis/03-0330HairSampleTesting-Community.pdf)

General Public



## Hair Sample Testing:

What can hair sampling results tell me about environmental exposures?

Sometimes health professionals can check whether people have been exposed to chemicals in the environment. For example, levels of chemicals can be measured in the soils, drinking water, and air where people live and work, and from these measures the amount of chemicals that people might routinely contact can be estimated. For some chemicals, urine and blood samples can be collected from people to look for evidence of actual exposure. In recent years, health professionals have debated what hair samples can tell us about environmental exposure. In this fact sheet, the Agency for Toxic Substances and Disease Registry (ATSDR), a public health agency of the U.S. Department of Health and Human

Services, looks at how best to interpret results from human hair samples to evaluate environmental exposures.

Currently, hair analysis is used for purposes other than assessing environmental exposures. For example, hair analysis has been used to test for illegal drug use and to conduct criminal investigations. This fact sheet does not address these other uses.

### Summary

ATSDR believes many scientific issues need to be resolved before hair analysis can become a useful tool to understand environmental exposures. Although hair analysis may answer some questions about environmental exposure to a few substances, hair analysis often raises more questions than it answers.

### What are hair samples and hair analyses?

A *hair sample* is a collection of hair strands, which are most commonly cut from a person's head. Hair samples usually contain hair that has grown over the last 12 months. Although hair growth varies, 12 months of hair growth usually represents about 5 inches of hair. *Hair analysis* occurs when laboratories measure the amount of specific substances in the hair sample.

Unfortunately, no widely accepted standards specify how hair samples should be collected, stored, and analyzed, and different laboratories use different methods when conduct-

ing hair analysis. Therefore, it is possible that two laboratories will report different results for hair samples collected from the same person. The different approaches used to collect and analyze hair samples make it very difficult for health professionals to know what the results of any individual hair sample really means.

### If a substance is detected in my hair, does that mean I have been exposed to environmental contamination?

Not necessarily. Generally, substances can end up in your hair in two ways. First, chemicals already in your body may get into your hair. These chemicals might have been in something you ate, the water you drank, or the air you breathed. Alternatively, the source of the chemicals could be from other exposures, or they could simply occur naturally in your body. Second, chemicals outside your body, like those in dust particles and hair care products, might stick to your hair after coming into contact with it.

Given how often most people cut their hair, hair analysis generally will not tell you anything about exposures that occurred more than one year ago. Furthermore, for most chemicals, hair analysis cannot tell you where a chemical in your hair came from. More specifically, when a chemical is detected in your hair, we often cannot tell if it is from a contaminated waste site, from your diet, or from other sources. This makes it very difficult to understand what a positive test result for a chemical in your hair truly means.

### Do health professionals commonly use hair analysis to diagnose health problems?

No. For most environmental contaminants, we simply do not know enough right now to use someone's hair analysis results to predict if health problems will occur. In fact, for many chemicals, we do not even know the range of levels that are typically found in the hair of an unexposed person. Without this information, we cannot tell if one person's hair

analysis result is unusually high or low. Because of these information gaps, doctors and other health professionals *rarely* use hair analysis to evaluate health problems. It is possible that future research will help us better understand what hair analysis results mean. Until this research is done, however, hair analysis results (with few exceptions) will not provide useful information about possible health problems.

### **With these limitations, why do health agencies and other groups collect hair samples?**

Many public health agencies, including ATSDR, have collected and analyzed hair samples when addressing community concerns regarding environmental exposure. Before deciding to use hair analysis, ATSDR carefully considers if the results can be interpreted and will add to our understanding of potential community exposure. In most cases, we choose not to collect hair samples because the results would not help us address health concerns. Given the limitations of hair analysis, health agencies almost never base health conclusions entirely on hair analysis results. Rather, hair analysis results are viewed as just one small piece of evidence that are considered along with other information when evaluating a site.

### **What is the bottom line? What can hair analysis results tell me?**

If an environmental chemical is detected in your hair, the only defensible conclusion you can draw, with few exceptions, is that you were exposed to the substance at some point over the last year or so. You will generally not know the source of the exposure or when it occurred. Most importantly, a detection will not tell you if your exposures might cause health problems.

*Remember:* For almost every environmental contaminant, hair analysis results alone will not tell you if you are likely to get sick or will have health problems.

### **Where can I get more information on how hair analysis relates to environmental exposure?**

ATSDR recently gathered a group of experts to discuss what scientists currently know and do not know about hair analysis. You can get a copy of the report that summarizes this expert panel meeting by calling ATSDR's toll-free telephone number: 1-888-42-ATSDR (or 1-888-422-8737). If you have access to the Internet, the report is available on ATSDR's Web site at "[www.atsdr.cdc.gov/HAC/hair\\_analysis](http://www.atsdr.cdc.gov/HAC/hair_analysis)." We can also send you a more detailed and technical fact sheet that summarizes our opinions on the current state of the science regarding hair analysis.

**For more information,  
contact ATSDR's toll-free  
information line:**

**(888) 42-ATSDR. . .  
that's (888) 422-8737**

**ATSDR's Internet address is [http://  
www.atsdr.cdc.gov](http://www.atsdr.cdc.gov)**